

# Comprehensive Opioid Abuse Site-Based Program (COAP) FY 2018

**Peter Ryba, PharmD**

PMP Director

[peter.ryba@state.nm.us](mailto:peter.ryba@state.nm.us)

**Maria Gonzales**

PMP Manager

[maria.gonzales@state.nm.us](mailto:maria.gonzales@state.nm.us)



## *COAP FY 2018 Competitive Grant Announcement*

---

- The purpose of the Harold Rogers PDMP is to improve collaboration and strategic decision-making among regulatory and law enforcement agencies and public health entities to address prescription drug and opioid abuse, save lives, and reduce crime.
- Announced May 8, 2018; due June 7, 2018

## *COAP FY 2018 Competitive Grant Announcement*

---

Enhancements Requested:

- [nmpmp.org](http://nmpmp.org) Updates (\$25,000)
- NarxCare (\$196,000/yr x2)
- PMP Integration (\$100,000)

Total (including required training, travel): \$527,800



## *NarxCare*

---

- Provides a more comprehensive approach to addressing substance use disorder.
- Aggregates and analyzes prescription information from providers and pharmacies and presents interactive, visual representations of that information.

## *NarxCare*

---

- Facilitates access to treatment through medication assisted treatment (MAT) locations,
- Facilitates patient education and engagement through CDC information sheets, and
- Enables the coordination of care via a communication module.



### NarxCare


- The Narx Report
- Resources
- Care Team Communications



### Narx Report

The Narx Report includes a patient's:

- Narx Scores
- Overdose Risk Scores
- Red Flags
- Rx Graph
- PMP Data






### Narx Report

#### Narx Score

- Type-specific use scores for opioids, sedatives, and stimulants
- Scores are quantified representations of the data in the PMP and range from 000-999.






### Narx Report

#### Narx Score

- Higher scores equate to higher risk and misuse
- The last digit represents the number of active prescriptions





## *Narx Report*

---

### Narx Score

- Algorithm incorporates the number of prescribers, number of pharmacies, milligram equivalents, and overlapping prescriptions
- Time element such that more recent activity is weighted more heavily
- Key requirement for overlapping prescriptions is that two different prescribers prescribe the same type of medication for use on the same day



## *Narx Report*

---

Distribution of Narx Scores:


- 75% score less than 200
- 5% score above 500
- 1% score above 650



### Narx Report

#### Red Flags


- Fully configurable and customizable PMP and non-PMP red flags
- When present, a red flag(s) may contribute to the risk of unintentional overdose or other adverse events.



### Narx Report

#### Overdose Risk Score

- Composite risk score which incorporates relevant data (PMP and non-PMP) into advanced customized predictive models to calculate overall risk.





## *Narx Report*

---

Trigger a discussion, not a decision

- Discuss risk/benefit of using a controlled substance.
- Discuss concerns with patient such as patterns for unsafe conditions, high doses with recent abstinence, multiple providers/pharmacies
- Consider naloxone prescription
- Consider a lock-in program
- Consider taper to lower dose or discontinuation
- Consider referral to MAT location

### Narx Report

#### Rx Graph

- Interactive display
- User can click into specific data points or click and drag to see additional detail
- Easily visualize and understand patterns in prescribing and usage behaviors






PRESCRIPTION MONITORING PROGRAM  
NEW MEXICO BOARD OF PHARMACY

Narx Report

PMP Data

- Displays full prescription detail as seen currently in PMP AWA<sub>RxE</sub>



### Resources

The NarxCare Resource module includes:

- MAT Locator
- CDC Resources

This module is fully configurable, so the State can add/remove forms at any point.

age: 30M date: 1/19/2017 | NARX REPORT | RESOURCES

**SAMHSA Buprenorphine Treatment Locator**

More information [here](#)

Step 1 Enter the zip code to center the search around

Step 2 Choose the radius for the search ☒ 10 mi ☐ 25 mi ☐ 50 mi

Step 3 Click submit and print form

First	Last	Dep.	Address	City	Cour
Dr. David	Beane	M.D.	1338 College Drive Suite B	Maricopa	West Court
Dr. Rakesh Kumar	Kaneria	M.D.	7760 West VOA Park Drive Suite G	West Chester	Butte Court
Dr. J.	Stafford	M.D.	1061 Bernard Road	New Vienna	Clint Court
Dr. Ramalingam	Selvarajah	M.D.	1649 Brice	Reynoldsburg	Franklin Court

**CDC Resources**

Click the associated link and print More information [here](#)

#### What You Need to Know

**PRESCRIPTION OPIOIDS: WHAT YOU NEED TO KNOW**

Prescription opioids can be used to help manage pain, but they can also be misused. This infographic provides information on how to use prescription opioids safely and what to do if you suspect someone is misusing them.

**WHAT ARE THE RISKS AND SIGNS OF OPIOID USE?**

Prescription opioids can be addictive. If someone is addicted, they may have withdrawal symptoms when they stop taking them. Signs of addiction include:

- Taking more than prescribed
- Taking them more often than prescribed
- Taking them for longer than prescribed
- Taking them for reasons other than pain
- Taking them with alcohol or other drugs
- Taking them when not in pain
- Taking them when not prescribed
- Taking them when not needed
- Taking them when not safe
- Taking them when not legal
- Taking them when not ethical
- Taking them when not responsible
- Taking them when not caring
- Taking them when not loving
- Taking them when not kind
- Taking them when not fair
- Taking them when not honest
- Taking them when not brave
- Taking them when not strong
- Taking them when not smart
- Taking them when not good
- Taking them when not great
- Taking them when not perfect
- Taking them when not happy
- Taking them when not healthy
- Taking them when not whole
- Taking them when not complete
- Taking them when not perfect
- Taking them when not happy
- Taking them when not healthy
- Taking them when not whole
- Taking them when not complete

**WHAT ARE THE RISKS AND SIGNS OF OPIOID USE?**

Prescription opioids can be addictive. If someone is addicted, they may have withdrawal symptoms when they stop taking them. Signs of addiction include:

- Taking more than prescribed
- Taking them more often than prescribed
- Taking them for longer than prescribed
- Taking them for reasons other than pain
- Taking them with alcohol or other drugs
- Taking them when not in pain
- Taking them when not prescribed
- Taking them when not needed
- Taking them when not safe
- Taking them when not legal
- Taking them when not ethical
- Taking them when not responsible
- Taking them when not caring
- Taking them when not kind
- Taking them when not fair
- Taking them when not honest
- Taking them when not brave
- Taking them when not strong
- Taking them when not smart
- Taking them when not good
- Taking them when not great
- Taking them when not perfect
- Taking them when not happy
- Taking them when not healthy
- Taking them when not whole
- Taking them when not complete
- Taking them when not perfect
- Taking them when not happy
- Taking them when not healthy
- Taking them when not whole
- Taking them when not complete

#### Opioids and Chronic Pain

**PROMOTING SAFER AND MORE EFFECTIVE PAIN MANAGEMENT**

Chronic pain is a complex condition that can be difficult to manage. This infographic provides information on how to manage chronic pain safely and effectively.

**SAFER AND MORE EFFECTIVE PAIN MANAGEMENT**

Chronic pain is a complex condition that can be difficult to manage. This infographic provides information on how to manage chronic pain safely and effectively.

**SAFER AND MORE EFFECTIVE PAIN MANAGEMENT**

Chronic pain is a complex condition that can be difficult to manage. This infographic provides information on how to manage chronic pain safely and effectively.

#### Pregnancy and Opioids

**PREGNANCY AND OPIOID PAIN MEDICATIONS**

Women who take opioid pain medications during pregnancy may be at risk of complications for themselves and their babies. This infographic provides information on how to manage pain safely during pregnancy.

**PREGNANCY AND OPIOID PAIN MEDICATIONS**

Women who take opioid pain medications during pregnancy may be at risk of complications for themselves and their babies. This infographic provides information on how to manage pain safely during pregnancy.

**PREGNANCY AND OPIOID PAIN MEDICATIONS**

Women who take opioid pain medications during pregnancy may be at risk of complications for themselves and their babies. This infographic provides information on how to manage pain safely during pregnancy.



## *Care Team Communication*

---

- Enables communication
- Transmission of documents
- Will need to see what is customizable within this module; however, this module can be disabled for all users.

## *PMP Integration*

---

- On a national level, PMP have low provider use due to numerous access barriers, including the time required to access and obtain the PMP report (~4 minutes per request).
- The CDC reported examples of 7-, 80-, and 148-fold increases in the number of PMP reports used by clinicians after workflow integration.
- NM Department of Health hired a PMP-EHR Coordinator to facilitate conversations, and to fund integrations.



PRESCRIPTION MONITORING PROGRAM

NEW MEXICO BOARD OF PHARMACY

*Questions?*

---

